

ON ASYMPTOTIC GAUSS-LUCAS THEOREM

RIKARD BØGVAD, DMITRY KHAVINSON, AND BORIS SHAPIRO

ABSTRACT. In this note we extend the Gauss-Lucas theorem on the zeros of the derivative of a univariate polynomial to the case of sequences of univariate polynomials whose almost all zeros lie in a given convex bounded domain in \mathbb{C} .

1. INTRODUCTION

The celebrated Gauss-Lucas theorem claims that for any univariate polynomial $P(z)$ with complex coefficients, all roots of $P'(z)$ belong to the convex hull of the roots of $P(z)$, see Theorem 6.1 of [5]. Many generalizations have been obtained over the years see, e.g., [1, 2, 7] and references therein.

In the present note, motivated by problems in potential theory in \mathbb{C} , we extend the Gauss-Lucas theorem to sequences of polynomials of increasing degrees whose almost all zeros lie in a given convex bounded domain in \mathbb{C} . Namely, given a convex bounded domain $\Omega \subset \mathbb{C}$, let $\{p_n(z)\}_{n=0}^\infty$ be a sequence of univariate polynomials with the degrees $\deg p_n = m_n$ such that $\lim_{n \rightarrow \infty} m_n = +\infty$. Assume that $\lim_{n \rightarrow \infty} \frac{\#_n(\Omega)}{m_n} = 1$, where $\#_n(\Omega)$ is the number of zeros of p_n lying in Ω (counted with multiplicities).

Problem 1. *Following the above notation we now ask whether there exists $\lim_{n \rightarrow \infty} \frac{\#'_n(\Omega)}{m_n - 1}$, where $\#'_n(\Omega)$ denotes the number of zeros of $p'_n(z)$ lying in Ω ?*

It turns out that the answer to Problem 1 formulated verbatim as above, is, in general, negative.

Example 1. Let O be the open square $(-2, 2) \times (-4i, 0)$. If $T_n(z) := \cos(n \arccos z)$ is the n -th Chebyshev polynomial of the first kind, then the derivative of $(z-i)T_n(z)$ has all its zeros in the upper half plane. Therefore, if we replace z by $z + ia_n$ for some sufficiently small a_n (i.e., shift all zeros downward by a_n), then we obtain polynomials of degrees $n + 1$ with n zeros in O , but whose derivatives have no zeros




FIGURE 1. Zeros of $((z - i)T_{30}(z))'$




FIGURE 2. Zeros of P (larger dots) and P' (smaller dots) for $P = (z^{10} - 1)(z - 3 + 2I)(z + 3 - 2I)(z - 1 + 5I)(z + 5)$ (left) and $P = Q(z - 3)(z + 2 - 2I)(z - 1 + I)(z + 2)$ where Q has 7 uniformly placed zeros on each side of the unit square (right).

in O . Choosing a_n appropriately for each n , we get a sequence of polynomials with all but one zeros in O whose derivatives have no zeros in O .

Strict convexity (e.g., as in the case of the open unit disk) will not be of much help either. Just replace above z by $M_n z$ with some large M_n and then make a vertical translation so that after all these operations the image of $[-1, 1]$ becomes a tiny secant segment of the unit circle. (This example was suggested to the third author by Professor V. Totik.)

However with slightly weaker requirements Problem 1 has a positive answer.

Theorem 1. *Given a polynomial sequence $\{p_n(z)\}$ as above and any $\epsilon > 0$,*

$$\lim_{n \rightarrow \infty} \frac{\#'_n(\Omega_\epsilon)}{m_n - 1} = 1,$$

where $\#'_n(\Omega_\epsilon)$ is the number of zeros of $p'_n(z)$ lying in Ω_ϵ , and Ω_ϵ is the open ϵ -neighborhood of $\Omega \subset \mathbb{C}$.

Two illustrations of Theorem 1 are given in Figure 2.

Acknowledgements. The third author want to thank Professor V. Totik for discussions of the problem.

2. PROOF

We first prove Theorem 1 in the case when Ω is a disk. Fix $\epsilon > 0$. Let $\mathbb{D} = \{|z| < 1\}$ be the open unit disk, $p_n(z) = \prod_{k=1}^{m_n} (z - \alpha_k)$, $\lim m_n = \infty$. Let us factor p_n as follows:

$$(1) \quad p_n = q_n r_n = \prod_{k=1}^{k_n} (z - a_k) \prod_{k=k_n+1}^{m_n} (z - b_k), \quad |b_k| > 1 + \epsilon, \quad |a_k| < 1 + \epsilon, \quad \text{with} \quad \lim_{n \rightarrow \infty} \frac{k_n}{m_n} = 1.$$

Denote by $\mathfrak{Z}'_n := \#\{z : |z| < 1 + \epsilon : p'_n(z) = 0\}$. We want to show that $\lim_{n \rightarrow \infty} \frac{\mathfrak{Z}'_n}{m_n - 1} = 1$.

Let

$$(2) \quad \hat{\mu}_n(z) := \frac{1}{m_n} \sum_{k=1}^n \frac{1}{z - \alpha_k} = \frac{1}{m_n} \frac{p'_n}{p_n},$$

denote the Cauchy transform of the root-counting probability measure μ_n of p_n . Note that

$$(3) \quad \hat{\mu}_n(z) = \frac{1}{m_n} \left(\frac{q'_n}{q_n} + \frac{r'_n}{r_n} \right) = \frac{1}{m_n} (k_n \hat{\nu}_n + (m_n - k_n) \hat{\psi}_n),$$

where ν_n and ψ_n are the root-counting measures of q_n and r_n respectively. All zeros of q'_n lie in the unit disk D by the Gauss-Lucas theorem. Also (1) implies that

$$(4) \quad \frac{(m_n - k_n)}{m_n} \|\psi_n\| \rightarrow 0.$$

Formula (4) implies that for all $p : 1 \leq p < 2$, we have

$$(5) \quad \left\| \frac{1}{m_n} \frac{r'_n}{r_n} \right\|_{L^p(dA)} \rightarrow 0$$

on compact subsets of \mathbb{C} . Here, $dA = \frac{1}{\pi} dx dy$ denote the normalized area measure.

Equation (5) follows from a trivial observation. Let μ be a Borel measure with a compact support. Then for any compact set $K \subset \mathbb{C}$, and any $p : 1 \leq p < 2$, we have

$$(6) \quad \|\hat{\mu}(z)\|_{L^p(K, dA)}^p \leq C(p, K) \|\mu\|.$$

Indeed, $\hat{\mu} = \int \frac{d\mu(\xi)}{\xi - z}$, hence

$$(7) \quad \int_K |\hat{\mu}|^p dA \leq \|\mu\| \int_{|\xi| < R} \frac{1}{|\xi|^p} dA \leq C \|\mu\|,$$

where R is chosen so that $\forall \xi \in K$ the disk of radius R centered at ξ contains K . The integral in (7) converges for all $p < 2$ and (6) follows, hence does (5).

Thus, we have from (5) the following corollary.

Corollary 1. . For any fixed $R > 1 + \epsilon$, and any $p : 1 \leq p < 2$, for almost all $r : 1 + \epsilon < r < R$, we have

$$(8) \quad \lim_{n \rightarrow \infty} \frac{1}{m_n} \int_{|z|=r} \frac{|r'_n|^p}{|r_n|^p} ds_r = 0,$$

where ds_r is the arclength measure on $\{z : |z| = r\}$.

Thus, from (3), we now obtain

Corollary 2.

$$\lim_{n \rightarrow \infty} \frac{1}{m_n} \int_{|z|=r} \left| \frac{p'_n}{p_n} - \frac{q'_n}{q_n} \right|^p ds_r = 0$$

for almost all $r : 1 + \epsilon < r < R$ and $p : 1 \leq p < 2$.

However

$$\frac{1}{m_n} \left(\frac{p'_n}{p_n} - \frac{q'_n}{q_n} \right) = \frac{1}{m_n} \left(\sum_{k=k_n+1}^{m_n} \frac{1}{z - b_k} \right)$$

by (3), and hence is analytic inside $\{|z| < 1 + \epsilon\}$ since $|b_k| > 1 + \epsilon$.

Therefore, from standard results on Hardy spaces H^p in the disk, cf. [3], we conclude that

$$(9) \quad \frac{1}{m_n} \left(\frac{p'_n}{p_n} - \frac{q'_n}{q_n} \right) \rightarrow 0$$

uniformly in the closed disk $\bar{D} = \{|z| \leq 1 + \epsilon\}$.

Since $\frac{1}{m_n} \frac{q'_n}{q_n}$ vanishes at $k_n - 1$ points in D by Gauss-Lucas theorem, invoking Hurwitz's theorem, we conclude that

$$(10) \quad \lim_{n \rightarrow \infty} \frac{1}{m_n} [\#(z : |z| < 1 + \epsilon : p'_n(z) = 0) - (k_n - 1)] = 0.$$

Since, by assumption, $\lim_{n \rightarrow \infty} \frac{k_n - 1}{m_n} = \lim_{n \rightarrow \infty} \frac{k_n}{m_n} = 1$, we arrive at

$$\lim_{n \rightarrow \infty} \frac{\mathfrak{Z}_n'}{m_n} = \lim_{n \rightarrow \infty} \frac{\mathfrak{Z}_n'}{m_n - 1} = 1,$$

which settles Theorem 1 in the case of a disk.

To finish the proof for the general case of an arbitrary convex domain Ω observe that we only used some properties of a disk to get a convenient foliation of a neighborhood of the unit disk by concentric circles and when applying Gauss-Lucas theorem. Both these facts are readily available for an arbitrary bounded convex domain. Finally, the Hardy spaces of analytic functions in the disk are replaced by the Smirnov classes E^p of functions representable by Cauchy integrals with L^p -densities (with respect to arclength). In the domains with piecewise smooth boundaries, e.g., convex domains, the latter behave in the very same manner as Hardy spaces – cf. [3]. \square

REFERENCES

- [1] J. S-C. Cheng, On the Distribution of the Critical Points of a Polynomial, preprint, June 2012.
- [2] B. Čurgus and V. Mascioni, A contraction of the Lucas polygon, Proc. of AMS, Volume 132, Number 10, Pages 2973–2981.
- [3] P. L. Duren, Theory of H^p Spaces, Academic Press, New York, 1970.
- [4] J. Garnett, Analytic Capacity and Measure. Lect. Notes Math. Vol. 297, Springer-Verlag, iv+138 pp., 1972.
- [5] M. Marden, Geometry of Polynomials, AMS, 1966.
- [6] Q. I. Rahman und G. Schmeisser, Analytic Theory of Polynomials, Oxford University Press, 2002.
- [7] A. Rüdinger, Strengthening the Gauss-Lucas theorem for polynomials with zeros in the interior of the convex hull, arXiv:1405.0689v1.

DEPARTMENT OF MATHEMATICS, STOCKHOLM UNIVERSITY, S-10691, STOCKHOLM, SWEDEN
E-mail address: rikard@math.su.se

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTH FLORIDA, TAMPA, 33620, USA
E-mail address: dkhavins@usf.edu

DEPARTMENT OF MATHEMATICS, STOCKHOLM UNIVERSITY, S-10691, STOCKHOLM, SWEDEN
E-mail address: shapiro@math.su.se