

# ALGEBRAS OF CURVATURE FORMS ON HOMOGENEOUS MANIFOLDS

ALEXANDER POSTNIKOV, BORIS SHAPIRO, AND MIKHAIL SHAPIRO

To Dmitry Borisovich Fuchs with love from former students of Moscow 'Jewish' University

ABSTRACT. Let  $C(X)$  be the algebra generated by the curvature two-forms of standard holomorphic hermitian line bundles over the complex homogeneous manifold  $X = G/B$ . The cohomology ring of  $X$  is a quotient of  $C(X)$ . We calculate the Hilbert polynomial of this algebra. In particular, we show that the dimension of  $C(X)$  is equal to the number of independent subsets of roots in the corresponding root system. We also construct a more general algebra associated with a point on a Grassmannian. We calculate its Hilbert polynomial and present the algebra in terms of generators and relations.

## 1. HOMOGENEOUS MANIFOLDS

In this section we remind the reader the basic notions and notation related to homogeneous manifolds  $G/B$  and root systems, as well as fix our terminology.

Let  $G$  be a connected complex semisimple Lie group and  $B$  its Borel subgroup. The quotient space  $X = G/B$  is then a compact homogeneous complex manifold. We choose a maximal compact subgroup  $K$  of  $G$  and denote by  $T = K \cap B$  its maximal torus. The group  $K$  acts transitively on  $X$ . Thus  $X$  can be identified with the quotient space  $K/T$ .

By  $\mathfrak{g}$  we denote the Lie algebra of  $G$  and by  $\mathfrak{h} \subset \mathfrak{g}$  its Cartan subalgebra. Also denote by  $\mathfrak{g}_{\mathbb{R}} \subset \mathfrak{g}$  the real form of  $\mathfrak{g}$  such that  $i\mathfrak{g}_{\mathbb{R}}$  is the Lie algebra of  $K$ . Analogously,  $\mathfrak{h}_{\mathbb{R}} = \mathfrak{h} \cap \mathfrak{g}_{\mathbb{R}}$  and  $i\mathfrak{h}_{\mathbb{R}}$  is the Lie algebra of the maximal torus  $T$ . The *root system* associated with  $\mathfrak{g}$  is the set  $\Delta$  of nonzero vectors (roots)  $\alpha \in \mathfrak{h}^*$  for which the root spaces

$$\mathfrak{g}_{\alpha} = \{x \in \mathfrak{g} \mid [h, x] = \alpha(h)x \text{ for all } h \in \mathfrak{h}\}$$

are nontrivial. Then  $\mathfrak{g}$  decomposes into the direct sum of subspaces

$$\mathfrak{g} = \mathfrak{h} \oplus \sum_{\alpha \in \Delta} \mathfrak{g}_{\alpha}.$$

For  $\alpha \in \Delta$ , the spaces  $\mathfrak{g}_{\alpha}$  and  $\mathfrak{h}_{\alpha} = [\mathfrak{g}_{\alpha}, \mathfrak{g}_{-\alpha}]$  are one-dimensional and there exists a unique element  $h_{\alpha} \in \mathfrak{h}_{\alpha}$  such that  $\alpha(h_{\alpha}) = 2$ . The elements  $h_{\alpha} \in \mathfrak{h}$  are called *coroots*. Actually,  $\alpha \in \mathfrak{h}_{\mathbb{R}}^*$  and  $h_{\alpha} \in \mathfrak{h}_{\mathbb{R}}$ , for  $\alpha \in \Delta$ . Let us choose generators  $e_{\alpha} \in \mathfrak{g}_{\mathbb{R}}$  of the root spaces  $\mathfrak{g}_{\alpha}$  such that  $[e_{\alpha}, e_{-\alpha}] = h_{\alpha}$  for any root  $\alpha$ . Then  $[h_{\alpha}, e_{\alpha}] = 2e_{\alpha}$  and  $[h_{\alpha}, e_{-\alpha}] = -2e_{-\alpha}$ .

The root system  $\Delta$  is subdivided into a disjoint union of sets of positive roots  $\Delta_+$  and negative roots  $\Delta_- = -\Delta_+$  such that the direct sum  $\mathfrak{b} = \mathfrak{h} \oplus \sum_{\alpha \in \Delta_+} \mathfrak{g}_{\alpha}$  is the

---

*Date:* January 17, 1999.

Lie algebra of Borel subgroup  $B$ . The Weyl group  $W$  is the group generated by the reflections  $s_\alpha : \mathfrak{h}^* \rightarrow \mathfrak{h}^*$ ,  $\alpha \in \Delta_+$ , given by

$$s_\alpha : \lambda \mapsto \lambda - \lambda(h_\alpha) \alpha.$$

The lattice  $\hat{T} = \{\lambda \in \mathfrak{h}^* \mid \lambda(h_\alpha) \in \mathbb{Z} \text{ for all } \alpha \in \Delta\}$  is called the *weight lattice*. Every weight  $\lambda \in \hat{T}$  determines an irreducible unitary representation  $\pi_\lambda : T \rightarrow \mathbb{C}^\times$  of the maximal torus  $T$  given by  $\pi_\lambda(\exp(x)) = e^{\lambda(x)}$ , for  $x \in \mathfrak{h}_\mathbb{R}$ , and every irreducible unitary representation of  $T$  is of this form.

For a weight  $\lambda \in \hat{T}$ , the homomorphism  $\pi_\lambda : T \rightarrow \mathbb{C}^\times$  extends uniquely to a holomorphic homomorphism  $\bar{\pi}_\lambda : B \rightarrow \mathbb{C}^\times$ . Thus any  $\lambda \in \hat{T}$  determines a holomorphic line bundle  $L_\lambda = G \times_B \mathbb{C} = K \times_T \mathbb{C}$  over  $X = G/B = K/T$ . The line bundle  $L_\lambda$  has a canonical  $K$ -invariant hermitian metric.

The classical Borel's theorem [2] describes the cohomology ring  $H^*(X, \mathbb{C})$  of the homogeneous manifold  $X$  in terms of generators and relations:

$$H^*(X, \mathbb{C}) \cong \text{Sym}(\mathfrak{h}^*)/I_W,$$

where  $I_W$  is the ideal in the symmetric algebra  $\text{Sym}(\mathfrak{h}^*)$  generated by the  $W$ -invariant elements without constant term. The natural projection from  $\text{Sym}(\mathfrak{h}^*)$  to  $H^*(X, \mathbb{C})$  is the homomorphism that sends a weight  $\lambda \in \hat{T}$  to the first Chern class  $c_1(L_\lambda)$  of the line bundle  $L_\lambda$ .

The purpose of this article is to extend the cohomology ring  $H^*(X, \mathbb{C})$  to the level of differential forms on  $X$ . It is possible to exhibit differential two-forms that represent the Chern classes  $c_1(L_\lambda)$  in the de Rham cohomology of the homogeneous manifold  $X$ . Recall that for a holomorphic hermitian line bundle  $L : E \rightarrow X$  there is a canonically associated connection on  $E$ . Denote by  $\Theta(L)$  the *curvature form* of this connection, which is a differential two-form on  $X$ . Then the form  $i\Theta(L)/2\pi$  represents  $c_1(L)$ .

In order to construct the curvature forms  $\Theta(L_\lambda)$  explicitly, we define the elements  $e^\alpha \in \mathfrak{g}_\mathbb{R}^*$ ,  $\alpha \in \Delta$ , by  $e^\alpha(\mathfrak{h}) = 0$  and  $e^\alpha(e_\beta) = \delta_{\alpha\beta}$ , for any  $\beta \in \Delta$ . (Here  $\delta_{\alpha\beta}$  is Kroneker's delta.) The space of left  $K$ -invariant differential one-forms on  $K$  can be identified with the dual to its Lie algebra, t.e., with  $i\mathfrak{g}_\mathbb{R}^*$ . Thus the elements  $i e^\alpha$  can be regarded as one-forms on  $K$ . The differential two-form on  $K$  given by

$$\phi_\alpha = e^\alpha \wedge e^{-\alpha}, \quad \alpha \in \Delta,$$

is invariant with respect to the right translation action of the torus  $T$ . Thus  $\phi_\alpha$  produces a two-form on the manifold  $X$ , for which we will use the same notation  $\phi_\alpha$ . It is clear from the definition that  $\phi_{-\alpha} = -\phi_\alpha$ .

The following statement is implicit in [4].

**Proposition 1.** *For  $\lambda \in \hat{T}$ , the curvature form of the holomorphic hermitian line bundle  $L_\lambda$  is given by*

$$\Theta(L_\lambda) = \sum_{\alpha \in \Delta_+} \lambda(h_\alpha) \phi_\alpha.$$

Let  $\Phi$  be the algebra generated by the two-forms  $\phi_\alpha$ ,  $\alpha \in \Delta_+$ . The relations in  $\Phi$  are relatively simple:

$$\phi_\alpha \phi_\beta = \phi_\beta \phi_\alpha, \quad (\phi_\alpha)^2 = 0.$$

Thus  $\Phi$  is a  $2^N$ -dimensional algebra, where  $N = |\Delta_+|$ . The main object in this paper is the subalgebra of  $\Phi$  generated by the curvature forms  $\Theta(L_\lambda)$ .

## 2. MAIN RESULTS

Denote by  $C(X)$  the subalgebra in the algebra of differential forms on  $X$  that is generated by the curvature forms  $\Theta(L_\lambda)$  of line bundles. Obviously,  $C(X)$  has the structure of a graded ring:  $C(X) = C^0(X) \oplus C^1(X) \oplus C^2(X) \oplus \dots$ , where  $C^k(X)$  is the subspace of  $2k$ -forms in  $C(X)$ . In order to formulate our main results about  $C(X)$  we need some extra notation from the matroid theory.

Let  $V$  be a collection of vectors  $v_1, v_2, \dots, v_N$  in a vector space  $E$ , say, over  $\mathbb{C}$ . A subset of vectors in  $V$  is called *independent* if they are linearly independent in  $E$ . By convention, the empty subset is independent. Let  $\text{ind}(V)$  be number of all independent subsets in  $V$ .

A *cycle* is a minimal by inclusion not independent subset. For a cycle  $C = \{v_{i_1}, \dots, v_{i_l}\}$ , there is a unique, up to a factor, linear dependence  $a_1 v_{i_1} + \dots + a_l v_{i_l} = 0$  with non-zero  $a_i$ 's. Let us fix a linear order  $v_1 < v_2 < \dots < v_N$  of all elements of  $V$ . For an independent subset  $S$  in  $V$ , a vector  $v \in V \setminus S$ , is called *externally active* if the set  $S \cup \{v\}$  contains a cycle  $C$  and  $v$  is the minimal element of  $C$ . Let  $\text{act}(S)$  be the number of externally active vectors with respect to  $S$ .

**Theorem 2.** *The dimension of the algebra  $C(X)$  is equal to the number  $\text{ind}(\Delta_+)$  of independent subsets in the set of positive roots  $\Delta_+$ . Moreover, the dimension of the  $k$ -th component  $C^k(X)$  is equal to the number of independent subsets  $S \subset \Delta_+$  such that  $k = N - |S| - \text{act}(S)$ , where  $N = |\Delta_+|$ .*

We remark here that, although the number  $\text{act}(S)$  of externally active vectors depends upon a particular order of elements in  $V$ , the total number of subsets  $S$  with fixed  $|S| + \text{act}(S)$  does not depend upon a choice of ordering.

We will actually prove a more general result about an arbitrary collection of vectors  $V$ . Let  $V$  and  $E$  be as above. We will assume that the elements  $v_1, \dots, v_N$  of  $V$  span the  $n$ -dimensional space  $E$ . Thus  $N \geq n$ . Let  $F = \mathbb{C}^N$  be the linear space with a distinguished basis  $\phi^1, \dots, \phi^N$ . Then  $V$  defines the projection map  $p : F \rightarrow E$  that sends the  $i$ th basis element  $\phi^i$  to  $v_i$ . The dual map  $p^* : E^* \rightarrow F^*$  defines an  $n$ -dimensional plane  $P = \text{Im}(p^*)$  in  $F^*$ . In other words, the collection of vectors  $V$  can be identified with an element  $P$  of the Grassmannian  $G(n, N)$  of  $n$ -dimensional planes in  $\mathbb{C}^N$ .

Let  $\phi_1, \dots, \phi_N$  be the basis in  $F^*$  dual to the chosen basis in  $F$ . Denote by  $\Phi_N$  the quotient of the symmetric algebra  $\text{Sym}(F^*)$  modulo the relations  $(\phi_i)^2 = 0$ ,  $i = 1, \dots, N$ . Let  $\mathcal{C}_V$  be the subalgebra in  $\Phi_N$  generated by the elements of the  $n$ -dimensional plane  $P \subset F^*$ . In other words, the algebra  $\mathcal{C}_V$  is the image of the induced mapping

$$\text{Sym}(E^*) \longrightarrow \Phi_N = \text{Sym}(F^*) / \langle \phi_i^2, i = 1, \dots, N \rangle.$$

The algebra  $\mathcal{C}_V$  has an obvious grading  $\mathcal{C}_V = \mathcal{C}_V^0 \oplus \mathcal{C}_V^1 \oplus \mathcal{C}_V^2 \oplus \dots$  by degree of elements.

Suppose  $E = \mathfrak{h}$  and  $V$  is the collections of coroots  $h_\alpha$ ,  $\alpha \in \Delta_+$ . For  $\lambda \in \mathfrak{h}^* = E^*$ ,  $p^*(\lambda) = \Theta(L_\lambda) \in F^*$  is the curvature form (see Proposition 1). Then  $\mathcal{C}_V = C(X)$  is the algebra generated by the curvature forms  $\Theta(L_\lambda)$ .

In general, we have the following result.

**Theorem 3.** *The dimension of the algebra  $\mathcal{C}_V$  is equal to the number  $\text{ind}(V)$  of independent subsets in  $V$ . Moreover, the dimension of the  $k$ -th component  $\mathcal{C}_V^k$  is equal to the number of independent subsets  $S \subset V$  such that  $k = N - |S| - \text{act}(S)$ .*

We can also describe the algebra  $\mathcal{C}_V$  as a quotient of a polynomial ring. Let us say that a hyperplane  $H$  in  $E$  is a *V-essential hyperplane* if the elements of the subset  $\{v_i, i = 1, \dots, N \mid v_i \in H\}$  span the hyperplane  $H$ . Obviously, an essential hyperplane is uniquely determined by the subset of indices  $I_H = \{i \in \{1, \dots, N\} \mid v_i \notin H\}$ . We will call such subset  $I_H$  a *V-essential index subset*. Denote by  $d(H) = d_V(H) = |I_H|$  the number of its elements. A nonzero vector  $\lambda \in E^*$  determines the hyperplane  $H = \{x \in E \mid \lambda(x) = 0\}$  in  $E$ . Vectors  $\lambda_H \in E^*$  corresponding to essential hyperplanes  $H$  will be called *V-essential vectors*. They are defined up to a nonzero factor.

**Theorem 4.** *The algebra  $\mathcal{C}_V$  is naturally isomorphic to the quotient of the polynomial ring  $\text{Sym}(E^*)/\mathcal{I}_V$ , where the ideal  $\mathcal{I}_V$  is generated by the powers  $(\lambda_H)^{d(H)+1}$  of V-essential vectors for all V-essential hyperplanes  $H$  in  $E$ . The isomorphism is induced by the embedding  $p^* : E^* \rightarrow F^*$ .*

**Remark 5.** There are several equivalent definitions of essential subsets, as follows:

**1.** An index subset  $I = \{i_1, \dots, i_k\} \subset \{1, \dots, N\}$  is *V-essential* if and only if the following two conditions are satisfied: (i) the coordinate plane  $\langle \phi_{i_1}, \dots, \phi_{i_k} \rangle$  in  $F^*$  has one-dimensional intersection with the plane  $P$ ; (ii) there is no proper subset in  $I$  that satisfies the condition (i). For an V-essential hyperplane  $H$ , the vector  $p^*(\lambda_H) \in F^*$  spans the one-dimensional intersection of  $P$  and the coordinate plane associated with  $I_H$ .

**2.** Let  $\theta_1, \dots, \theta_n$  be any basis in  $P$ . The V-essential subsets are in one-to-one correspondence with the cycles in the vector set  $\{\phi_1, \dots, \phi_N, \theta_1, \dots, \theta_n\}$ . For a cycle  $\{\phi_{i_1}, \dots, \phi_{i_k}, \theta_{j_1}, \dots, \theta_{j_k}\}$ , the subset  $\{i_1, \dots, i_k\}$  is V-essential. Moreover, every V-essential subset is of this form.

Note that the decomposition of the Grassmannian  $G(n, N)$  of all  $n$ -dimensional planes  $P \subset F^*$  into strata with the same collection of essential subsets coincides with the decomposition of  $G(n, N)$  into small cells of Gelfand-Serganova [3] because any two  $P_1, P_2 \in G(n, N)$  with the same collection of essential subsets have the same dimensions of intersections with all coordinate subspaces. Equivalently,  $P_1$  and  $P_2$  are in the same strata if and only if the corresponding collections of vectors  $V_1$  and  $V_2$  define the same matroid, i.e., have the same collection of independent subsets.

Let us apply the Theorem 4 to  $C(X)$ . Let  $\omega_1, \omega_2, \dots, \omega_l$  be the fundamental weights. They generate the weight lattice  $\hat{T}$ . Also let  $d_i$  be the number of positive roots  $\alpha \in \Delta_+$  such that  $\alpha(\omega_i) \neq 0$ .

**Corollary 6.** *The algebra  $C(X)$  is naturally isomorphic to the quotient of the polynomial ring  $\text{Sym}(\mathfrak{h}^*)/\mathcal{J}$ , the ideal  $\mathcal{J} \in \text{Sym}(\mathfrak{h}^*)$  is generated by the elements  $(w \cdot \omega_i)^{d_i+1}$ , where  $i = 1, \dots, l$  and  $w$  is an element of the Weyl group  $W$ . This isomorphism is induced by the projection  $\text{Sym}(\mathfrak{h}^*) \rightarrow C(X)$  that send  $\lambda \in \hat{T}$  to the curvature form  $\Theta(L_\lambda)$ .*

For type  $A$  flag manifolds this statement was earlier proved in [7].

*Proof* — The generators of the ideal  $\mathcal{J}$ , as described in Theorem 4, correspond to root subsystems in  $\Delta$  of codimension 1. Every subsystem of codimension 1 is

congruent, modulo the Weyl group, to the root subsystem in the hyperplane  $H_i$  of zeroes of a fundamental weight  $\omega_i$ . Moreover,  $d(H_i) = d_i$  is exactly the number of positive roots that do not belong to this hyperplane.  $\square$

### 3. DUAL STATEMENT AND PROOF OF THEOREMS 2 AND 3

Next we give a dual version of Theorem 3. Let  $\text{Sym}(E)$  be the symmetric algebra of the vector space  $E$ . For a subset  $S \subset V$ , denote  $m(S) = \prod_{v \in S} v \in \text{Sym}(E)$ , a square-free monomial in  $v_i$ 's. Let us denote by  $\mathcal{S}_V$  a subspace in  $\text{Sym}(E)$  spanned by  $2^N$  square-free monomials  $m(S)$ , where  $S$  ranges over all subsets in  $V$ . Let  $\mathcal{S}_V^k$  be the  $k$ -th graded component of  $\mathcal{S}_V$ .

**Theorem 7.** *The dimension of the subspace  $\mathcal{S}_V$  in  $\text{Sym}(E)$  is equal to the number  $\text{ind}(V)$  of independent subsets in  $V$ . Moreover, the dimension of  $\mathcal{S}_V^k$  is equal to the number of independent subsets  $S \subset V$  such that  $k = N - |S| - \text{act}(S)$ .*

Let us choose a basis  $x^1, \dots, x^n$  in  $E$ . Let  $x_1, \dots, x_n$  be the dual basis in  $E^*$ , and let  $\theta_i = p^*(x_i)$ ,  $i = 1, \dots, n$ , be the corresponding basis in  $P$ .

For  $I = (1 \leq i_1 < i_2 < \dots < i_k \leq N)$  and  $J = (1 \leq j_1 \leq j_2 \leq \dots \leq j_k \leq n)$ , we denote by  $a_{IJ}$  the coefficient of the square-free monomial  $\phi_{i_1} \dots \phi_{i_k} \in \Phi_N$  in the product of generators  $\theta_{j_1} \dots \theta_{j_k} \in \mathcal{C}_V$ . Let  $A_k = (a_{IJ})$  be the  $\binom{N}{k} \times \binom{n+k-1}{k}$ -matrix formed by the  $a_{IJ}$ . Clearly,  $\dim \mathcal{C}_V^k$  is equal to the rank of the matrix  $A_k$ .

On the other hand,  $a_{IJ}$  is also the coefficient of the monomial  $x^{j_1} \dots x^{j_k} \in \text{Sym}(E)$  in the square-free monomial  $v_{i_1} \dots v_{i_k} \in \mathcal{S}(V)$ . Thus  $\dim \mathcal{S}_V^k$  is equal to the rank of the transposed matrix  $A_k^T$ , which is the same as the rank of  $A_k$ . We proved the following statement, which implies the equivalence of Theorems 3 and 7.

**Lemma 8.** *For  $k = 1, \dots, N$ , we have  $\dim \mathcal{C}_V^k = \dim \mathcal{S}_V^k$ .*

Let us say that a subset  $S \subset V$  is *robust* if there is no cycle  $C \subset V$  with minimal element  $v$  such that  $S \cap C = \{v\}$ .

**Lemma 9.** *The number of robust subsets in  $V$  is equal to the number of independent subsets. Moreover, the number of  $k$ -element robust subsets is equal to the number of independent subsets  $S$  with  $k = N - |S| - \text{act}(S)$ .*

*Proof* — We present an explicit bijection between robust subsets and independent subsets. A subset  $A$  is the complement to a robust subset in  $V$  if and only if for any cycle  $C$  with minimal element  $v$ , inclusion  $C \setminus \{v\} \subset A$  implies  $C \subset A$ . We will call such subsets *antirobust*.

For an independent subset  $S$ , let  $M$  be the collection of all externally active  $v \in V \setminus S$ . Then  $S \cup M$  is antirobust. Conversely, for an antirobust subset  $A$ , let  $M$  be the collection of all  $v \in V$  such that  $v$  is a minimal element in some cycle  $C \subset A$ . Then  $A \setminus M$  is an independent subset.

Clearly, both these mappings are inverse to each other and the statement of lemma follows.  $\square$

**Theorem 10.** *The set of square-free monomials  $m(S)$ , where  $S$  ranges over robust subsets, forms a basis of the subspace  $\mathcal{S}_V$ .*

First, we prove a weaker version of Theorem 10.

**Lemma 11.** *The square-free monomials  $m(S)$ , where  $S$  ranges over robust subsets, span  $\mathcal{S}_V$ .*

*Proof* — Suppose not. Let  $m(R)$  be the maximal in the lexicographical order square-free monomial which cannot be expressed linearly via the monomials  $m(S)$  with robust  $S$ . Then there is a cycle  $C = \{v_{i_1}, v_{i_2}, \dots, v_{i_l}\}$  with the minimal element  $v = v_{i_1}$  such that  $R \cap C = \{v\}$ . We can replace  $v$  in the monomial  $m(R)$  by a linear combination of  $v_{i_2}, v_{i_3}, \dots, v_{i_l}$ . Thus  $m(R)$  is a linear combination of square-free monomials which are greater than  $m(R)$  in the lexicographical order. By assumption each of these monomials can be expressed via the monomials  $m(S)$  with robust  $S$ . Contradiction.  $\square$

We can now conclude the proof.

*Proof of Theorems 7 and 10* — Lemmas 9 and 11 imply the inequality  $\dim \mathcal{S}_V \leq \text{ind}(V)$ . In view of these two lemmas it is enough to show that  $\dim \mathcal{S}_V$  is actually equal to  $\text{ind}(V)$ .

We prove this statement by induction on  $|V|$ . If the linear span of vectors in  $V$  is one-dimensional, then both  $\dim \mathcal{S}_V$  and  $\text{ind}(V)$  are equal to the number of non-zero vectors in  $V$  plus one. This establishes the base of induction.

Assume that  $v = v_N$  is a nonzero vector in  $V$ . Let  $V' = V \setminus \{v\} = \{v_1, \dots, v_{N-1}\}$ , and let  $V''$  be the collection of images of vectors  $v_1, \dots, v_{N-1}$  in the quotient space  $E/\langle v \rangle$ . It follows from the definition of independent subset that  $\text{ind}(V) = \text{ind}(V') + \text{ind}(V'')$ . Assume by induction that  $\dim \mathcal{S}_{V'} = \text{ind}(V')$  and  $\dim \mathcal{S}_{V''} = \text{ind}(V'')$ .

Clearly,  $\mathcal{S}_V$  is spanned by  $\mathcal{S}_{V'} + \mathcal{S}_{V'}v$ . Both  $\mathcal{S}_{V'}$  and  $\mathcal{S}_{V'}v$  have same dimensions. Hence,  $\dim \mathcal{S}_V = 2 \dim \mathcal{S}_{V'} - \dim(\mathcal{S}_{V'} \cap \mathcal{S}_{V'}v)$ . Let  $\pi : \mathcal{S}_{V'} \rightarrow \mathcal{S}_{V''}$  be the natural projection. Then  $\mathcal{S}_{V'} \cap \mathcal{S}_{V'}v \subset \text{Ker}(\pi)$ . Thus  $\dim \mathcal{S}_{V'} - \dim(\mathcal{S}_{V'} \cap \mathcal{S}_{V'}v) \geq \dim \mathcal{S}_{V''}$  and  $\dim \mathcal{S}_V \geq \dim \mathcal{S}_{V'} + \dim \mathcal{S}_{V''} = \text{ind}(V') + \text{ind}(V'') = \text{ind}(V)$ . Coupled with the inequality  $\dim \mathcal{S}_V \leq \text{ind}(V)$ , this produces the required statement.

This finishes the proof of Theorems 7 and 10 and thus of Theorems 2 and 3.  $\square$

#### 4. PROOF OF THEOREM 4

Let  $\text{Ess}_V$  denote the set of all  $V$ -essential hyperplanes in  $E$ . Recall that  $\mathcal{I}_V$  is the ideal in  $\text{Sym}(E^*)$  generated by the powers of essential vectors  $(\lambda_H)^{d(H)+1}$ ,  $H \in \text{Ess}_V$  (see Theorem 4). The embedding  $p^* : E^* \rightarrow F^*$  induces the mapping  $\text{Sym}(E^*) \rightarrow \Phi_N$ , whose image is the algebra  $\mathcal{C}_V$ . Let  $\tilde{\mathcal{I}}_V$  denote the kernel of this mapping, which we will call the *vanishing ideal* of  $V$ . Theorem 4 amounts to the identity of ideals  $\mathcal{I}_V = \tilde{\mathcal{I}}_V$ .

The proof relies on a couple of simple lemmas. As in the previous section we assume that  $v_N$  is a nonzero vector in  $V$ . Let  $V' = V \setminus \{v_N\} = \{v_1, \dots, v_{N-1}\}$ , and let  $V''$  be the collection of images of vectors  $v_1, \dots, v_{N-1}$  in the quotient space  $E/\langle v_N \rangle$ . We also denote by  $\text{Ess}_{V'}$  and  $\text{Ess}_{V''}$  the sets of  $V'$ -essential and  $V''$ -essential hyperplanes in the corresponding spaces.

The dimension of the span of vectors in  $V''$  is  $n - 1$ . The dimension of the span of  $V'$  can be either  $n - 1$  or  $n$ . For a hyperplane  $H$  in  $E/\langle v_N \rangle$ , let  $\bar{H} = H \oplus \langle v_N \rangle$  be the hyperplane in  $E$ . Also for a collection of hyperplanes  $C$  in  $E/\langle v_N \rangle$ , let  $\bar{C} = \{\bar{H} \mid H \in C\}$  be the collection of hyperplanes in  $E$ .

**Lemma 12.** (a) *If  $\dim V' = n - 1$  then  $\text{Ess}_V = \{\langle V' \rangle\} \cup \overline{\text{Ess}_{V''}}$ .*

(b) If  $\dim V' = n$  then  $\text{Ess}_V = \text{Ess}_{V'} \cup \overline{\text{Ess}_{V''}}$ . For  $H \in (\text{Ess}_{V'} \setminus \overline{\text{Ess}_{V''}})$ , we have  $d_V(H) = d_{V'}(H) + 1$ . For  $H \in \overline{\text{Ess}_{V''}}$ , we have  $d_V(H) = d_{V''}(H)$ .

*Proof* — The part (a) is left as an easy exercise for the reader. In order to prove (b) we first assume that a hyperplane  $H$  contains  $v_N$ . Then  $H$  is  $V$ -essential if and only if its projection  $H/\langle v_N \rangle$  is  $V''$ -essential. Suppose that a hyperplane  $H$  does not contain  $v_N$ . Then  $H$  is  $V$ -essential if and only if it is  $V'$ -essential and its projection  $H/\langle v_N \rangle$  is not  $V''$ -essential. The equalities for the numbers  $d(H)$  are also obvious. This proves the lemma.  $\square$

Recall that the collection of vectors  $V$  is associated with a plane  $P \in G(n, N)$ . Let  $P'$  and  $P''$  be the planes associated with vector sets  $V'$  and  $V''$ , respectively. Namely,  $P'$  is the projection of  $P$  along  $\phi_N$  onto the hyperplane  $\mathcal{H} = \langle \phi_1, \dots, \phi_{N-1} \rangle$  spanned by the first  $N - 1$  coordinate vectors; and  $P''$  is the intersection of  $P$  with the same hyperplane  $\mathcal{H}$ .

We can choose the basis  $x_1, \dots, x_n$  in  $E^*$  and the corresponding basis  $\theta_i = p^*(x_i)$ ,  $i = 1, \dots, n$  in  $P$  such that  $\theta_1, \dots, \theta_{n-1} \in \mathcal{H}$  and  $\theta_n \in \phi_N + \mathcal{H}$ . Also denote  $\tilde{\theta}_n = \theta_n - \phi_N \in \mathcal{H}$ . Then  $\theta_1, \dots, \theta_{n-1}, \tilde{\theta}_n$  span the projected space  $P'$ . The space  $\text{Sym}(E^*)$  can be identified with the polynomial ring  $\mathbb{C}[x_1, \dots, x_n]$

**Lemma 13.** *The vanishing ideal  $\tilde{\mathcal{I}}_V$  consists of all polynomials  $f \in \mathbb{C}[x_1, \dots, x_n]$  such that both  $f$  and  $\partial f / \partial x_n$  belong to the vanishing ideal  $\tilde{\mathcal{I}}_{V'}$ . In particular,  $\tilde{\mathcal{I}}_V \subseteq \tilde{\mathcal{I}}_{V'}$ .*

*Proof* — Recall that the ideal  $\tilde{\mathcal{I}}_V$  consists of all polynomials in  $x_1, \dots, x_n$  which vanish in the algebra  $\Phi_N$  upon substituting of the  $\theta_i$  instead of the  $x_i$ . Taylor's expansion in the algebra  $\Phi_N$  gives

$$f(\theta_1, \dots, \theta_n) = f(\theta_1, \dots, \tilde{\theta}_n + \phi_N) = f(\theta_1, \dots, \tilde{\theta}_n) + (\partial f / \partial x_n)(\theta_1, \dots, \theta_{n-1}, \tilde{\theta}_n) \phi_N.$$

We have only two non-vanishing terms in the right hand side, since  $\phi_N^2 = 0$ .

The polynomial  $f(x_1, \dots, x_n)$  belongs to  $\tilde{\mathcal{I}}_V$  if and only if both  $f(\theta_1, \dots, \tilde{\theta}_n)$  and  $(\partial f / \partial x_n)(\theta_1, \dots, \theta_{n-1}, \tilde{\theta}_n)$  vanish in the algebra  $\Phi_N$ .  $\square$

We now conclude the proof of the identity  $\mathcal{I}_V = \tilde{\mathcal{I}}_V$ . The inclusion  $\mathcal{I}_V \subseteq \tilde{\mathcal{I}}_V$  is straightforward. Indeed, every  $p^*(\lambda_H)$  is a linear combination of  $d(H)$  different  $\phi_i$ 's. Thus  $(\lambda_H)^{d(H)+1}$  maps to zero in the algebra  $\Phi_N$ .

We prove the identity  $\mathcal{I}_V = \tilde{\mathcal{I}}_V$  by induction on  $|V|$ . The base of induction is the trivial case  $V = \{v_1\}$ . For  $|V| \geq 2$ , assume by induction that the statement is true for both  $V'$  and  $V''$ . Take any  $f \in \tilde{\mathcal{I}}_V$ . Our goal is to show that  $f \in \mathcal{I}_V$ .

By Lemma 13, the polynomial  $\partial f / \partial x_n$  belongs to  $\tilde{\mathcal{I}}_{V'}$ . Notice that for any  $H \in \text{Ess}_{V'} \cap \overline{\text{Ess}_{V''}}$  the coordinate expansion of the corresponding  $V'$ -essential vector  $\lambda_H$  does not involve  $x_n$ . Indeed, any  $H \in \overline{\text{Ess}_{V''}}$  contains  $v_N$ , thus  $\lambda_H(v_N) = 0$ . On the other hand,  $x_1(v_N) = \dots = x_{n-1}(v_N) = 0$ , and  $x_n(v_N) = 1$ .

By inductive assumption, one has

$$\partial f / \partial x_n = \sum_{H \in \text{Ess}_{V'} \setminus \overline{\text{Ess}_{V''}}} p_H \lambda_H^{d_{V'}(H)+1} + \sum_{H \in \text{Ess}_{V'} \cap \overline{\text{Ess}_{V''}}} p_H \lambda_H^{d_{V'}(H)+1},$$

where the  $p_H$  are certain polynomials in  $x_1, \dots, x_n$ . The  $\lambda_H$  in the second sum do not involve  $x_n$ . Thus, integrating the above expression with respect to  $x_n$ , one

deduces that there exists a polynomial  $\bar{f}$  of the form

$$\bar{f} = \sum_{H \in \text{Ess}_{V'} \setminus \overline{\text{Ess}_{V''}}} \bar{p}_H \lambda_H^{d_{V'}(H)+2} + \sum_{H \in \text{Ess}_{V'} \cap \overline{\text{Ess}_{V''}}} \bar{p}_H \lambda_H^{d_{V'}(H)+1}$$

satisfying  $\partial \bar{f} / \partial x_n = \partial f / \partial x_n$ . By Lemma 12, the polynomial  $\bar{f}$  can be written as

$$\bar{f} = \sum_{H \in \text{Ess}_V} \bar{p}_H \lambda_H^{d_V(H)+1}$$

and thus belongs to  $\mathcal{I}_P$ . The difference  $\hat{f} = f - \bar{f}$  belongs to  $\tilde{\mathcal{I}}_P$  and is independent on  $x_n$ . Thus  $\hat{f} \in \tilde{\mathcal{I}}_{V''}$ . By induction hypothesis,  $\hat{f} \in \mathcal{I}_{V''} \subset \mathcal{I}_V$ . Thus  $f = \bar{f} + \hat{f} \in \mathcal{I}_V$ . The statement follows. Q.E.D.

## 5. REMARKS AND OPEN PROBLEMS

The algebra  $C(X)$  for type A flag manifolds  $X = SL(n)/B$  was studied in more details in [7] and [6] motivated by [1]. In this case, we first proved Theorem 2 in [6] using a different approach based on a presentation of  $C(X)$  as a quotient of a polynomial ring (cf. Theorem 4). The theorem claims that the dimension of  $C(X)$  is equal to the number forests on  $n$  labelled vertices whereas the dimension of  $C^k(X)$  is equal to the number of forests with  $\binom{n}{2} - k$  inversions. The statement about forests was initially conjectured in [7] and the statement concerning inversions was then guessed by R. Stanley. In [6] we also discuss various generalizations of the ring  $C(X)$ .

A natural open problem is to extend the results to homogeneous manifolds  $G/P$ , where  $P$  is a parabolic subgroup. Formulas for curvature forms on  $G/P$  can be found in [4].

It is also intriguing to find the links between the algebra  $C(X)$  and the arithmetic Schubert calculus, see H. Tamvakis [8, 9].

B. Kostant pointed out that the algebra  $C(X)$  is related to the  $S(\mathfrak{g})$ -module  $\wedge \mathfrak{g}$  studied in [5]. It would be interesting to investigate this relationship.

The authors are grateful to Vladimir Igorevich Arnold, Richard Stanley, Alek Vainshtein, and Andrei Zelevinsky for stimulating discussions and helpful suggestions.

## REFERENCES

- [1] V. I. Arnold, Remarks on eigenvalues and eigenvectors of Hermitian matrices, Berry phase, adiabatic connections and quantum Hall effect *Selecta Math.* **(1)** (1995), 1–19.
- [2] A. Borel, Sur la cohomologie de espaces fibrés principaux et des espaces homogènes des groupes de Lie compacts, *Ann. of Math.* (2) **57** (1953), 115–207.
- [3] I. Gelfand and V. Serganova, Combinatorial geometries and the strata of a torus on homogeneous compact manifolds, *Russ. Math. Surv.* **42**, no. 2, (1987), 107–134.
- [4] P. Griffiths and W. Schmid, Locally homogeneous complex manifolds, *Acta Math.* **123** (1969), 253–302.
- [5] B. Kostant, On  $\wedge \mathfrak{g}$  for a semisimple Lie algebra  $\mathfrak{g}$ , as an equivariant module over the symmetric algebra  $S(\mathfrak{g})$ , preprint.
- [6] A. Postnikov, B. Shapiro, and M. Shapiro, Chern forms on flag manifolds and forests, in preparation.
- [7] B. Shapiro and M. Shapiro, On algebra generated by Bott-Chern 2-forms on  $SL_n/B$ , *C. R. Acad. Sci. Paris Sér. I Math.* **326** ser. 1 (1998), 75–80.
- [8] H. Tamvakis, Bott-Chern forms and arithmetic intersections, *Enseign. Math.* **43** (1997), 33–54.

- [9] H. Tamvakis, Arithmetic intersection theory on flag varieties, preprint alg-geom/9611006, 1996.

DEPT. MATH., MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139, U.S.A.  
*E-mail address:* `apost@math.mit.edu`

DEPT. MATH., UNIVERSITY OF STOCKHOLM, STOCKHOLM, S-10691, SWEDEN  
*E-mail address:* `shapiro@matematik.su.se`

DEPT. MATH., ROYAL INSTITUTE OF TECHNOLOGY, STOCKHOLM, S-10044, SWEDEN  
*E-mail address:* `mshapiro@math.kth.se`