

# On root asymptotics for the eigenpolynomials of a Bochner-Krall operator

joint with G. Masson, H. Rullgård, R. Bøgvad, J. Borcea,  
T. Bergkvist

Boris Shapiro

November 25, 2013

## Topics to discuss

- 1 Bochner-Krall problem, formulation and results
- 2 Bochner-Krall operators
  - Non-degenerate case
  - Degenerate case
- 3 Homogenized spectral problem
- 4 Heine-Stieltjes theory

## Main references

G. Masson and B. Shapiro, **On polynomial eigenfunctions of a hypergeometric-type operator**, *Exper. Math.*, vol. 10, 609-618, (2001).

T. Berqkvist and H. Rullgård, **On polynomial eigenfunctions for a class of differential operators**, *Math. Res. Lett.*, vol.9, 153-171, (2002).

T. Berqkvist, H. Rullgård and B. Shapiro, **On Bochner-Krall orthogonal polynomial systems**, *Math. Scand.*, vol 94, 148–154, (2004).

## Main references continued

J. Borcea, B. Shapiro, **Root asymptotics of spectral polynomials for the Lamé operator**, Comm.Math.Phys, vol 282 (2008) 323–337.

J. Borcea, R. Bøgvad and B. Shapiro, **Homogenized spectral pencils for exactly solvable operators: asymptotics of polynomial eigenfunctions**, Publ. RIMS, vol 45 (2009) 525–568.

T. Bergkvist, **On asymptotics of polynomial eigenfunctions for exactly-solvable differential operators**, J. Approx. Theory 149 (2007), no. 2, 151–187.

# Bochner-Krall problem

**Definition 1.** An operator  $T = \sum_{i=1}^k Q_i(z) \frac{d^i}{dz^i}$  is called **Bochner-Krall** if  $\deg Q_i(z) \leq i$  and there exists a value  $i$  such that  $\deg Q_i(z) = i$ .

Obviously,  $T(z^j) = a_j z^j + \text{lower order terms}$ , i.e.  $T$  acts diagonally in the monomial basis.

**Lemma.** For any Bochner-Krall operator  $T$  and sufficiently large  $n$  there exists a unique eigenpolynomial  $p_n(z)$  of degree  $n$ .

In fact, S.Bochner has shown in 1929 that any linear differential univariate operator having an infinite sequence of polynomial eigenfunctions is a Bochner-Krall operator.

## Bochner-Krall problem, cont.

### Problem (Bochner (1929), Krall (1938))

*Which Bochner-Krall operators have orthogonal polynomials (with respect to a positive or a signed measure supported on  $\mathbb{R}$ ) as their sequences of eigenpolynomials?*

### Lemma (Bochner)

*A Bochner-Krall operator having a sequence of orthogonal polynomials (BKOPS) must be of even order and formally self-adjoint.*

## Bochner-Krall problem, cont.

### Theorem (Bochner)

*The only second order BKOPS correspond to four classical orthogonal polynomial families: Jacobi, Bessel, Laguerre and Hermite polynomials.*

### Theorem (Krall)

*The fourth order BKOPS additionally contain three more families: the Legendre-type, the Laguerre-type, the Jacobi-type polynomials.*

## Bochner-Krall problem, cont.

For orders exceeding 4 the complete answer is still unknown!!!

All known BKOPS have distributional weights of the form  $w = u + v$ , where  $u$  is a classical orthogonal weight and  $v$  consists of some point masses supported on the boundary of the support of  $u$ .

It is conjectured that this is true for all BKOPS.

It is also conjectured that the leading coefficient of a BKOPS is a power of either linear or quadratic polynomial.



## Bochner-Krall problem, cont.

### Theorem (Kwon-Lee)

*The only BKOPS with compactly supported positive measure on  $\mathbb{R}$  as the Jacobi-type polynomials, i.e. after an appropriate linear real change of variables they are orthogonal w.r.t*

$$w = (1 - x)^\alpha (1 + x)^\beta H(1 - x^2) + c\delta(x - 1) + d\delta(x + 1),$$

*where  $\alpha, \beta > -1$  and  $c, d \geq 0$ , and  $H(x)$  is the Heaviside step function.*

# Main problem and examples

**Our main problem.** Given an arbitrary Bochner-Krall operator  $T$  describe the root asymptotics for the polynomial sequence  $\{p_n(z)\}$ .

## Examples

$$T_1 = z(z-1)(z-1) \frac{d^3}{dz^3}$$

$$T_2 = (z-1)(z+1)(z-2+31)(z-3-21) \frac{d^4}{dz^4}$$

$$T_3 = (z-1)(z+1)(z-2+31)(z-3-21)(z+3) \frac{d^5}{dz^5}$$

$$T_4 = (z^2+1)(z-2+31)(z-3-21)(z+3)(z+1+1) \frac{d^6}{dz^6}$$

# Roots of eigenpolynomials




Figure : Roots of  $p_{55}(z)$  for the above  $T$ 's.

## Basic Definitions

**Definition 3.** Cauchy transform of a (complex-valued) measure  $\rho$  satisfying  $\int_{\mathbb{C}} d\rho(\xi) < \infty$  is given by

$$C_{\rho}(z) = \int_{\mathbb{C}} \frac{d\rho(\xi)}{z - \xi}.$$

**Example.** If  $d\rho(z) = \frac{1}{\pi\sqrt{1-x^2}}$ ,  $x \in [-1, 1]$  then  $C_{\rho} = \frac{1}{\sqrt{z^2-1}}$  in  $\mathbb{C} \setminus [-1, 1]$

**Definition 1.1.** A Bochner-Krall operator  $T = \sum_{i=1}^k Q_i(z) \frac{d^i}{dz^i}$  is called of **non-degenerate** if  $\deg Q_k(z) = k$ .

## First results

**Proposition 1.2.** Assuming that each  $p_n(z)$  is monic and that  $\Psi(z) = \lim_{n \rightarrow \infty} \frac{p'_n(z)}{np_n(z)}$  exists in some open neighborhood  $\Omega$  of  $\mathbb{C}$  one gets that  $\Psi(z)$  satisfies in  $\Omega$  the algebraic equation

$$Q_k(z)\Psi^k(z) = 1.$$

**Theorem 1.1,** H. Rullgård. Let  $Q_k(z)$  be a monic degree  $k$  polynomial.  $\exists!$  probability measure  $\mu_Q$  such that

- supp  $\mu_Q$  is compact;
- its Cauchy transform  $C_\mu$  satisfies the equation  $Q_k(z)C_\mu^k(z) = 1$  almost everywhere in  $\mathbb{C}$ .

# Main Theorem

**Main theorem.** In the above notation

1)  $\text{supp } \mu_Q$  is a curvilinear tree which is straightened out by the analytic mapping

$$\xi(z) = \int_a^z \frac{dz}{\sqrt[k]{Q_k(z)}}.$$

2)  $\text{supp } \mu_Q$  contains all the zeros of  $Q_k(z)$  and is contained in the convex hull of those.

3) There is a natural formula for the angles between the branches and the masses of the branches satisfy Kirchhoff law.

# Illustration




Figure : The measure  $\mu_Q$  before and after the transformation.

Here  $Q(z) = (z - 1)(z - 3)(z - 1)$

## Illustration cont.




Figure : Example of a  $\mu_Q$  with angles.

**Speculation.** Distribution should be related to the algebraic curve  $y^k = Q_k(z)$  and Stokes lines for the differential operator.



## Degenerate case

A Bochner-Krall operator  $T$  of order  $k$  is called **degenerate** iff  $\deg Q_k < k$ .

Examples:  $T = z \frac{d^2}{dz^2} + (az + b) \frac{d}{dz}$ ,  $T = \frac{d^2}{dz^2} + (az + b) \frac{d}{dz}$   
leading to Laguerre resp. Hermite polynomials.

**Proposition.** The union of all roots of all polynomial eigenfunctions of a Bochner-Krall operator  $T$  is unbounded if and only if  $T$  is degenerate.

**Question.** Given a degenerate  $T$  with the family of eigenpolynomials  $\{p_n(z)\}$  how fast does the maximum modulus among the roots of  $p_n(z)$  grow?

# Main Conjecture

**Conjecture.** (T.Bergkvist) Given a degenerate  $T = \sum_{j=1}^k Q_j(z) \frac{d^j}{dz^j}$  denote by  $j_0$  the largest  $j$  for which  $\deg Q_j(z) = j$ . Then

$$\lim_{n \rightarrow \infty} \frac{r_n}{n^d} = c_T$$

where  $c_T > 0$  is a positive constant and

$$d := \max_{j \in [j_0+1, k]} \left( \frac{j - j_0}{j - \deg Q_j} \right).$$

# Main Conjectural Corollary

The Cauchy transform  $C(z)$  of the asymptotic root measure  $\mu$  of the scaled eigenpolynomial  $q_n(z) = p_n(n^d z)$  of a degenerate  $T$  satisfies the following algebraic equation for almost all complex  $z$ :

$$z^{j_0} C^{j_0}(z) + \sum_{j \in A} \alpha_{j, \deg Q_j} z^{\deg Q_j} C^j(z) = 1,$$

where  $A$  is the set consisting of all  $j$  for which the maximum  $d := \max_{j \in [j_0+1, k]} \left( \frac{j-j_0}{j-\deg Q_j} \right)$  is attained, i.e.  
 $A = \{j : (j - j_0)/(j - \deg Q_j) = d\}$ .

# More Pictures




Figure : Degenerate case.

# What do we need to solve Bochner-Krall problem?

## Problem (Seems doable)

*Assuming that Bergkvist's conjecture is settled, deduce the information about the support and the density of the measure from the above algebraic equation.*

In particular,

## Conjecture

*The support of the measure satisfying a.e. the algebraic equation of T.Bergkvist is always a tree.*

# What do we need to solve Bochner-Krall problem? cont.

## Problem

*Assuming that  $\{p_n(x)\}$  is an OPS with respect to a positive measure with unbounded (connected?) support on  $\mathbb{R}$ , is there an asymptotic root-counting measure for the scaled roots of  $\{p_n(x)\}$ ?*

## Problem

*If such a measure exists, can one get it explicitly for some natural classes of OPS?*

(Such results are known if the support is compact but in the unbounded case there seems to be no information!)

## Homogenized stuff (non-degenerate case so far)

Only the leading coefficient was important in the previous set-up which is unfair! To improve the situation we use (following Wasow, Fedoryuk etc) the *homogenized spectral problem* of the form

$$T_\lambda = \sum_{i=0}^k Q_i(z) \lambda^{k-i} \frac{d^i}{dz^i},$$

where each  $Q_i(x) = a_{ij}z^j + a_{i,j-1}z^{j-1} + \dots$  is a polynomial of degree  $i$ .

## basic facts

**Definition.** A non-degenerate  $T$  is called **of general type** iff  $\sum_{i=0}^k a_{ij} \lambda^{k-i} = 0$  has  $k$  distinct zeros.

**Proposition 2.1.** If  $T$  is of general type,

- 1) for all sufficiently large  $n$  there exist exactly  $k$  distinct values  $\lambda_{n,j}$ ,  $j = 1, \dots, k$  of the spectral parameter  $\lambda$  such that the operator  $T_\lambda$  has a polynomial eigenfunction  $p_{n,j}(z)$  of degree  $n$ .
- 2) Asymptotically  $\lambda_{n,j} \sim n\lambda_j$  where  $\lambda_1, \dots, \lambda_k$  is the set of roots of the algebraic equation  $\sum_{i=0}^k a_{i,j} x^{k-i} = 0$ .




## basic conjectures

**Conjecture 1.** If  $T$  is of general type and all  $\lambda_1, \dots, \lambda_k$  have distinct arguments then for each  $j = 1, \dots, k$   $\exists!$  probability measure  $\mu_j$  with compact support whose Cauchy transform  $C_j(z)$  satisfies almost everywhere in  $\mathbb{C}$

$$\sum_{i=1}^k Q_i(z) (\lambda_j C_j(z))^i = 0.$$

**Conjecture 2.**  $C_j(z) = \lim_{n \rightarrow \infty} \frac{p'_{n,j}(z)}{\lambda_{n,j} p_{n,j}(z)}$  outside the support of  $\mu_j$  which is the union of finitely many segments of analytic curves.

# Even more nice pictures



## Remarks

**Observation.** Near  $\infty \in \mathbb{CP}^1$  the Cauchy transforms  $\lambda_1 G_1(z), \dots, \lambda_k G_k(z)$  are independent sections of the symbol equation of  $T_\lambda$  considered as a branched cover over  $\mathbb{CP}^1$ .

**Open Problems.** Find "explicit" description of the measures  $\mu_i$  and their supports. What is their relation to the plane curve  $\sum_{i=1}^k Q_i(z)y^i = 0$ ?

# Heine-Stieltjes theory

Take a general linear operator  $T = \sum_{i=0}^k Q_i(z) \frac{d^i}{dz^i}$  with polynomial coefficients and set

$$r = \max_i (\deg Q_i(z) - i).$$

If  $r \geq 0$ ,  $\deg Q_k(z) = k + r$  and  $Q_k(z)$  has at least two distinct roots we call  $T$  **general Lamé-type** operator.

Consider the generalized spectral problem

$$T(p(z)) + V(z)p(z) = 0,$$

where  $p(z)$  is an eigenpolynomial and  $V(z)$  is a spectral polynomial. (Classically,  $p(z)$  is called a **Stieltjes** polynomial and  $V(z)$  is called a **Van Vleck** polynomial.) Note that  $\deg V(x) \leq r$ .

# Results

**Proposition 3.1.** Under the above assumptions for any sufficiently large  $n$  there exist exactly  $\binom{n+r}{r}$  degree  $n$  Stieltjes polynomials  $p_{n,j}(z)$  and corresponding Van Vleck polynomials  $V_{n,j}(z)$ .

## Results, cont.

**Proposition 3.2.** If a sequence  $\{\tilde{V}_{n,j_n}(z)\}$ ,  $n = 1, \dots$ , of scaled Van Vleck polynomials converge to some polynomial  $\tilde{V}(z)$  then the sequence of finite measures  $\mu_{n,j}$  of the corresponding family of eigenpolynomials  $\{p_{n,j_n}(z)\}$  converge to a measure  $\mu_{\tilde{V}}$  satisfying the properties:

- supp  $\mu_{\tilde{V}}$  is a forest of curvilinear trees;
- the union of the leaves of supp  $\mu_{\tilde{V}}$  coincides with the union of all zeros of  $Q_k(z)$  and that of  $\tilde{V}(z)$ .
- supp  $\mu_{\tilde{V}}$  is straightened out by the transformation given by

$$\int_a^z \frac{\tilde{V}(z) dz}{Q_k(z)}.$$

## Example

$$T = (z^2 + 1)(z + 2I - 3)(z - 3I - 2) \frac{d^3}{dz^3}$$




Figure : Examples of  $\mu_Q$ 's for the above  $T$ .

## Example, cont.




Figure : Union of  $\mu_Q$ 's for the above  $T$ .



## Example, cont.





Figure : The union of roots for the Van Vleck polynomials for the above  $T$ .

# Problems



**Figure :** The union of roots for the Van Vleck polynomials for  $T = Q_5(z) \frac{d^4}{dz^4}$ .

The asymptotic distributions for the roots of Van Vleck polynomials are completely unknown for operators of order  $> 2$ !

# Thanks for your patience!